


# Risque relatif de thrombose veineuse et contraception oestro progestative selon une cohorte danoise de 8 millions de femmes

## Type de contraception orale



Progestogen with 50 µg ethinylestradiol:	
Norethisterone	
Levonorgestrel	
Progestogen with 30-40 µg ethinylestradiol:	
Norethisterone	
Phasic levonorgestrel	
Levonorgestrel combined	
Norgestimate	
Desogestrel	
Gestodene	
Drospirenone	
Cyproterone	
Progestogen with 20 µg ethinylestradiol:	
Desogestrel	
Gestodene	
Drospirenone	



## Risque relatif

5.66 (3.12 to 10.3)	5,6
3.54 (2.48 to 5.05)	
1.57 (0.84 to 2.92)	1,7
2.28 (1.85 to 2.83)	
2.19 (1.74 to 2.75)	
2.56 (2.18 to 3.01)	
4.21 (3.63 to 4.87)	
4.23 (3.87 to 4.63)	
4.47 (3.91 to 5.11)	
4.10 (3.37 to 4.99)	
3.26 (2.88 to 3.69)	
3.50 (3.09 to 3.97)	4,8
4.84 (3.19 to 7.33)	

## Sans Contraception orale



## Risque relatif

1

*Risk of venous thromboembolism from use of oral contraceptives containing different progestogens and oestrogen doses:  
Danish cohort study, 2001-9. BMJ 2011;343:d6423 doi*